



HB1550 | 115 RM

GDU to Mid-Silk N/A
 GDU to Black Layer N/A
 Pollination for Maturity N/A

Hybrid Highlights

- Good overall agronomics
- Impressive performance against key traited checks
- Best performance in zone with good southern movement as well
- Strong overall disease package, including *Goss's wilt*
- Dual-purpose potential

Agronomics

Staygreen	Above Avg
Greensnap	Above Avg
Stalks	Very Good
Roots	Above Avg
Early Vigor	Very Good
Drought Tolerance	Very Good
Test Weight	Very Good
Silage	Very Good

Water Management

Full Irrigation	HR
Limited Irrigation	HR
Rainfed	HR
Dryland (Stress)	HR

Disease Tolerance

N. Corn Leaf Blight	Very Good
Gray Leaf Spot	Very Good
Southern Leaf Blight	Very Good
Goss's Wilt	Very Good
Common Rust	Above Avg
Southern Rust	N/A
Stalk Rot	Above Avg
Ear Rot	Above Avg

Management Response

Added Management	Very Good
Fungicide Response	Above Avg
Average Management	Very Good
Low Management	Very Good

Trait: CONV

Region Adaptability




Soil Placement

Course (Droughty)	Very Good
Medium	Excellent
Heavy (Well Drained)	Very Good
Heavy (Poorly Drained)	Above Avg
Variable	Excellent

Rotation Management

Rotated Acres	HR
Continuous Corn	HR
Continuous Corn w/ Fungicide	R

Key





HB1550 | 115 RM

HARVEST BOUNTY® CORN

Plant Description

Plant Height	Med-Tall
Ear Height	Med-High
Leaf Angle	Semi-Upright
Leaf Color	N/A
Leaf Width	N/A
Silk Color	N/A
Anther Color	N/A
Kernel Cap Color	N/A
Kernel Row	18-20
Cob Color	Pink
Ear Length	Semi-Long
Ear Girth	Girthy
Ear Type	Semi-Flex
Husk Cover	Medium

Population Management

Yield Environment	Population Range
0-100	14000-20000
101-150	20000-28000
151-200	28000-32000
201-250	32000-35000
251-300	35000-42000

Herbicide Sensitivity

Growth Regulator	Acceptable
Sulfonylureas Inhibitors (ALS)	Acceptable
Pigment Inhibitors (HPPD)	Acceptable

Yield Environment Placement

Tough	Excellent
Variable	Excellent
High Yield	Very Good



Performance may vary from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and consider the impacts of these conditions on the grower's fields. Always follow grain marketing and IRM requirements and pesticide label directions. Agronomic characteristics and ratings may vary with growing conditions and environment. Ratings are approximate and should not be considered as absolute. Ratings on new products are based on limited data and may change as more data are collected. Extreme or variable conditions may adversely affect performance. WILBUR-ELLIS logo and HARVEST BOUNTY are registered trademarks, HARVEST BOUNTY logo is a trademark of Wilbur-Ellis Company LLC.