



HB1690W | 116 RM

GDU to Mid-Silk N/A
 GDU to Black Layer N/A
 Pollination for Maturity Med-Late

Hybrid Highlights

- Attractive, food-grade white corn hybrid with very good overall agronomics
- Strong stalks with impressive late-season intactness
- Strong resistance to ear molds, specifically Fusarium ear molds
- Widely adapted east to west
- Dual-purpose potential

Agronomics

Staygreen	Above Avg
Greensnap	Above Avg
Stalks	Very Good
Roots	Very Good
Early Vigor	Above Avg
Drought Tolerance	Above Avg
Test Weight	Excellent
Silage	Above Avg

Water Management

Full Irrigation	HR
Limited Irrigation	HR
Rainfed	HR
Dryland (Stress)	HR

Disease Tolerance


N. Corn Leaf Blight	Above Avg
Gray Leaf Spot	Very Good
Southern Leaf Blight	N/A
Goss's Wilt	N/A
Common Rust	Above Avg
Southern Rust	N/A
Stalk Rot	Very Good
Ear Rot	Very Good

Management Response

Added Management	Above Avg
Fungicide Response	Above Avg
Average Management	Excellent
Low Management	Excellent

Trait: CONV WHITE

Region Adaptability




Soil Placement

Course (Droughty)	Above Avg
Medium	Excellent
Heavy (Well Drained)	Very Good
Heavy (Poorly Drained)	Above Avg
Variable	Very Good

Rotation Management

Rotated Acres	HR
Continuous Corn	R
Continuous Corn w/ Fungicide	R

Key





HB1690W | 116 RM

HARVEST BOUNTY® CORN

Plant Description

Plant Height	Med-Tall
Ear Height	Med-High
Leaf Angle	Semi-Upright
Leaf Color	N/A
Leaf Width	N/A
Silk Color	N/A
Anther Color	N/A
Kernel Cap Color	N/A
Kernel Row	16-18
Cob Color	White
Ear Length	Average
Ear Girth	Average
Ear Type	Semi-Flex
Husk Cover	Good

Population Management

Yield Environment	Population Range
0-100	14000-20000
101-150	20000-28000
151-200	28000-32000
201-250	32000-35000
251-300	NR

Herbicide Sensitivity

Growth Regulator	Acceptable
Sulfonylureas Inhibitors (ALS)	Acceptable
Pigment Inhibitors (HPPD)	Acceptable

Yield Environment Placement

Tough	Excellent
Variable	Excellent
High Yield	Above Avg



Performance may vary from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and consider the impacts of these conditions on the grower's fields. Always follow grain marketing and IRM requirements and pesticide label directions. Agronomic characteristics and ratings may vary with growing conditions and environment. Ratings are approximate and should not be considered as absolute. Ratings on new products are based on limited data and may change as more data are collected. Extreme or variable conditions may adversely affect performance. WILBUR-ELLIS logo and HARVEST BOUNTY are registered trademarks, HARVEST BOUNTY logo is a trademark of Wilbur-Ellis Company LLC.